

NEW TOUGH ROUNDUP® GEL READY TO USE WEEDKILLER - 100ml

KILLS TOUGH WEEDS TO THE ROOTS

- Kills Tree Stumps
- Kills Blackberry, Ivy, Onion Weed and other deep rooted weeds.

JUST TOUCH THE WEED

Use weedkillers safely. Always read the label.

roundup-garden.co.nz

ROUNDUP® TOUGH GEL

ACTIVE CONSTITUENT:

72 g/L Glyphosate (present as the potassium salt). This product is a garden weed and stump killer for use on unwanted vegetation like blackberry, ivy and other difficult weeds. Roundup® TOUGH GEL prevents stump re-growth but does not accelerate stump degradation, this process continues at natural speed.



Convolvulus



Blackberry



Stump

HOW AND WHEN TO USE

- Remove cap and foil seal, then prime [first use] by clicking the blue button about 20 times until the gel appears at the top of the applicator.
- Difficult Weeds – Apply Roundup® TOUGH GEL when the weeds are green and fully grown (just before flowering).
- Starting with the largest leaves first, apply to every third leaf.
- First effects are seen on the leaves in 10–15 days. Root kill may take up to 28 days dependent on weed type and weather conditions.
- Cut Stumps - Apply to the area around the exposed bark of freshly cut stump with 1 click for a 2-4 cm diameter stump or more clicks in ratio for bigger stumps.
- Stumps may not show effects, but should not regrow the next season.
- After use, clean the gel from the blue applicator top by wiping on a weed leaf, before securely replacing the blue cap.

NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED. DO NOT APPLY ON OR AROUND FOOD OR ANIMAL FEED CROPS.

WHAT IS ROUNDUP® TOUGH GEL

Roundup® TOUGH GEL is a multi-purpose weedkiller that kills large and difficult to control weeds such as blackberry, wandering willie, onionweed, etc.

It is also suitable for application to freshly cut stumps of various woody weeds, including Ivy and trees.

It is broken down by soil micro-organisms to leave the soil ready for replanting.

WHAT TO EXPECT FROM ROUNDUP® TOUGH GEL

Roundup® TOUGH GEL will kill garden weeds with a single application when applied according to instructions for use.

The first effects may be seen on the leaves between 10-15 days, depending on the type of weed and the weather conditions. Roundup® TOUGH GEL will work slower in cool and dry conditions and on mature perennial weeds.

Root kill can take up to 28 days, depending on the type of weed and the weather conditions. On stumps the effect might only be visible the following season when no re-growth occurs.

HOW TO USE

Against difficult and large weeds, leaf application.

Apply Roundup® TOUGH GEL when the weeds are green and fully grown (just before flowering). Starting with the largest leaves first, apply to every third leaf.

Dose:

WEEDS TYPE	CLICKS	LEAF AREA TO TREAT
Tough Weeds such as: Blackberry, Onion Weed and other deep rooted weeds.	1 click	30 x 20 cm
Young trees and other woody.	1 click	30 x 20 cm

Wait 5 days before digging/soil disturbance or cultivation, so that Roundup® TOUGH GEL can move into the roots.

Re-application may be necessary, if re-growth appears the following year.

Stump control:

Cut the woody weed or sapling trees above soil level and apply Roundup® TOUGH GEL to the whole stump surface, particularly the around the edges and bark of the stump.

One click for a stump of 2-4 cm diameter, more clicks for bigger trees.

Best time to apply is during summer and autumn.

Examples of controlled species are: **Ivy, Poplar, Willow, Eucalypt and other invasive species.**

RESTRICTIONS:

Avoid contact with cultivated plants and desired vegetation. To avoid transfer to lawns and other desired vegetation do not walk into treated areas until the product has dried. Keep children and pets out of the treated area until the Roundup® has dried. Rain within 6 hours of application reduces effectiveness and repeat treatment may be necessary.

INSTRUCTIONS FOR USE

TO OPEN:

1. Squeeze the 2 blue cap lips at the side of the bottle and lift the cap.
2. Remove the foil seal.
3. Before first use, prime the GEL bottle by pressing the blue button around 20 times until GEL appears at the top of the bottle.
4. Push/click the blue button to release the gel.

HOW TO USE AS A LEAF APPLICATION:

5. Apply by wiping the leaf surface. There is no need to apply to the whole leaf surface.
6. Apply Roundup® TOUGH GEL according to the dose rate specified under 'HOW TO USE'.



AFTER USE:

Clean the gel from the blue applicator top by wiping on a weed leaf, before securely replacing the blue cap by aligning both blue side lips at the same time.

STORAGE AND DISPOSAL:

Store in the closed, original container in a cool, dry place out of reach of children. DO NOT store in direct sunlight. Dispose of empty container by wrapping in a paper, placing in a plastic bag and putting in the garbage. DO NOT store at low temperatures.

SAFETY DIRECTIONS:

Avoid contact with eyes and skin. Wash hands after use.

FOR EMERGENCIES ONLY CALL **NZ 0800 734 607**.

FIRST AID:

If poisoning occurs, contact a doctor or NZ Poisons Information Centre: **0800 764 766**.

Distributor: Tui Products Ltd, Truman Lane, Mt Maunganui, New Zealand Tel: 07 575 2160

® Registered Trademark of Monsanto Technology LLC,
Used under license by Monsanto Australia Limited.

HSNO approval Number: HSR101087

Refer to www.epa.govt.nz for controls.

PROTECT FROM FROST



Keep children and pets
away from treated areas
until gel has dried

